

# Brazilian plants with possible action on the central nervous system—A study of historical sources from the 16th to 19th century

Melina Giorgetti, Giuseppina Negri, Eliana Rodrigues\*

*CEBRID, Department of Psychobiology at UNIFESP, Sao Paulo, Brazil*

Received 21 April 2006; received in revised form 8 August 2006; accepted 8 August 2006

Available online 12 August 2006

## Abstract

Brazil is a country rich in biodiversity, endemism, and cultural diversity, inhabited by different types of population. European expeditions and the migratory processes that began in the 16th century greatly contributed both to cultural diversity and to Brazilian popular therapeutics, and produced the first records on medicinal plants in Brazil. This study comprises a bibliographical survey of historic books found in Sao Paulo libraries (16th through 19th centuries) on medicinal plants exerting effects on the central nervous system (CNS). Thirty-four plants native to Brazil were selected from the reading of the books. Of these 34 plants, 13 were also recorded in ethnopharmacological studies among modern Brazilian communities and 16 have been studied phytochemically. Only eight have been the object of pharmacological studies, six of these, recently, with a request for a patent. Results showed that most of the species recorded in this study have been reported as medicinal for centuries, but have never been the object of pharmacological investigation down to the present time. Such results provide ideas for a selection of these species as potentially bioactive to be included in future pharmacological studies.

© 2006 Elsevier Ireland Ltd. All rights reserved.

*Keywords:* Medicinal plants; Central nervous system; Ethnopharmacology; Historic literature; Popular knowledge; Ethnobotany

## 1. Introduction

Brazil is a country rich in diversity and endemism, with a territory that includes five main biomas: the Amazon rain forest, the Cerrado brushlands, the Mata Atlântica rain forest, the Pantanal wetlands, and the Caatinga semi-arid desert. Brazil is also rich in cultural diversity and is inhabited by several types of traditional communities and ethnic groups such as Indigenous ethnic groups, Quilombola communities, and other traditional populations (Caboclo/river dwellers, Caiçara fishermen, the Jangadeiro raftsmen, babaçu gatherers, and rubber tappers), the latter being the result of miscegenation between native Indian, European, and African elements. Descendants of Europeans and Asians settled in Brazil during colonization and migratory processes begun at the start of the 16th century. This miscegenation enhanced the culture and rituals of some Brazilian communities and ethnic groups, enriching them from the ethnobotanical point of view.

Little was known about Brazil at the time of the discoveries. The first Jesuits, scientists, explorers, and settlers to arrive in Brazil (such as Padre Anchieta, Guilherme Piso and Von Martius) reported on all the characteristics they could observe (Kury, 2001); these writings are today regarded as the first ethnobotanical records on Brazilian medicinal plants and their uses by different populations.

In time, this knowledge took different paths—in part passed on from one person to another as people moved on (migration, journeys, etc.) to be later re-transmitted, diffused and broadly utilized by several populations in all parts of the country. However, another part of the knowledge was not passed on: its use became less and less frequent over the course of time and, not having been awarded the attention it deserved by the population and scientific community, was wholly forgotten. Only a few written records survived in books from the past centuries. Some studies outside Brazil have focused at the history of plant use from an ethnopharmacological perspective (Fabre, 2003; Heinrich and Teoh, 2004; Kufer et al., 2005; Heinrich et al., 2006).

The main objective of this study is, from a bibliographical survey of historic Brazilian literature, to recover therapeutic uses of plants native to Brazil, going back over centuries, and, con-

\* Corresponding author at: Department of Psychobiology, UNIFESP, R. Botucatu, 862 Ed. Ciências Biológicas (1º Andar), CEP 04023-062 Sao Paulo, Brazil. Tel.: +55 11 21490155; fax: +55 11 50842793.

E-mail address: [elirodri@psicobio.epm.br](mailto:elirodri@psicobio.epm.br) (E. Rodrigues).

comitantly, to search for phytochemical and pharmacological investigations. The data obtained in this study may serve as a basis to select the species to be investigated in future pharmacological and phytochemical studies.

Therefore, the aim of the present study matches the recent concept for ethnopharmacology, which is an “Interdisciplinary study of the physiological actions of plant, animal and other substances used in indigenous medicines of past and present cultures” (International Society of Ethnopharmacology, 2005).

## 2. Methodology

The investigation focused on historic books on popular use of medicinal plants found in Sao Paulo institutions. Included were: the University of Sao Paulo libraries, the Botanical Institute of the State of Sao Paulo, the Padre Anchieta Museum, and private collections owned by researchers José Ribeiro do Valle and Elisaldo Luís de Araújo Carlini, with over 40 years of their academic life devoted to the study of medicinal plants. The search yielded six books of great interest to the project, published between the 16th and the 19th centuries. Reports on plant species with therapeutic uses that might affect the central nervous system (CNS) were obtained by reading the books. This article focuses on the CNS, based on the fact that 450 million people in all of the world experience mental, neurological, or behavioral problems at some time (WHO, 2006).

The geographical origin of the species selected was verified using the six-volume Pío Corrêa (1926); books by Lorenzi (1992, 1998, 2003), Lorenzi and Matos (2002), Missouri Botanical Garden Website (2006) and specific flora found in large herbaria.

Bibliographical surveys concerning ethnopharmacological studies were made to verify if species recorded in the present study have uses in modern Brazilian communities. It was searched in Pubmed and in UNIFESP collection. This research was carried out in June 2006.

Efforts were made to identify modern pharmacological studies that might partially validate some of the possible effects related to uses encountered in the historical sources. Bibliographical surveys were carried out in Pubmed (August 2005) and in all of the Annals of the Brazilian Symposium of Medicinal Plants (from 1968 until 2004), one of the most relevant sources for Brazil. In 2006 (May), a survey was also conducted in Dr. Duke’s Phytochemical and Ethnobotanical Database.

The bibliographical research on all of the chemical classes found in the plants selected was carried out in the Web of Science (July 2006) utilizing the scientific name of each plant. Results of the survey were evaluated to certify that the article contained the chemical constituents analyzed in order to discard false results. Information was classified in “phytochemical classes” present in each plant: the chemical constituent was not mentioned so that comparisons and results might be visualized more clearly.

Every attempt was made, also, to verify whether the species identified in this study are available on the market in Brazil as pharmaceutical products. ANVISA (National Agency for Sanitary Vigilance) – a regulatory agency for Brazilian medication with a complete list of these products – was con-

sulted (<http://www.anvisa.gov.br/> July 2005). DEF (Dicionário de Especialidades Farmaceuticas 2004/2005), an up-to-date listing of medication legally approved in Brazil was also consulted.

Finally, a survey verified the existence of any patent related to the species recorded in the present study. Sites for registry of patents in Brazil, the United States, and Europe, were consulted, namely, the Instituto Nacional da Propriedade Industrial—INPI (<http://www.inpi.gov.br/>), the United States Patent Office—USPTO (<http://www.uspto.gov>) and the European Patent Office—ESPACENET, 2005 (<http://www.ep.espacenet.com>). This survey was conducted during August 2005.

## 3. Results

Only a few books were found on the subject of medicinal plants in the institutions researched, written in former centuries and in good enough condition for reading. Books in a poor state of conservation were not included in this study, due to difficulties in interpreting data. The books on medicinal plants of greatest relevance belonged to private collections or museum libraries. All books utilized in this research are new editions, published in the 20th century, of original books.

The six books selected were from the 16th century (one book), the 17th century (one), the 18th century (one), and the 19th century (three):

- Book 1: Notícia do Brasil (1587). Sousa, G.S. Brasiliensia Documenta. Volume VII. Sao Paulo, 1974.
- Book 2: “Zoobiblion” (Livro de Animais do Brasil (1654). Wagens, Z. Brasiliensia Documenta. Volume IV. Sao Paulo, 1964.
- Book 3: Plantas Fluminenses Descritas por Frei Veloso (1741–1811). Veloso, J.M.C. Separata de Anais da Biblioteca Nacional Volume 96. Rio de Janeiro, 1976.
- Book 4: Plantas Medicinaias do Brasil (1801–1812). Gomes, B.A. Brasiliensia Documenta. Volume V. Sao Paulo, 1972.
- Book 5: Viagem pelo Brasil (1817–1820). Spix and Martius. Volume III. Editora Melhoramentos. Sao Paulo, 1976.
- Book 6: Natureza, Doenças, Medicina e Remédios dos Índios Brasileiros (1844). Martius. Companhia Editorial Nacional. Sao Paulo, 1939.

The books were read and 93 plants with effect or uses possibly related to the CNS recorded as reported by the respective authors. Uses most cited were: to combat fever (febrifuges—20.4%), against pain (analgesic—20.4%) and those that alter consciousness (that produce “drunkenness”, that produce “numbness”—16.1%).

*Bignonia chica* Humb. and Bonpl. used as blood depurative was also included. Even though this category of use has no direct relationship with the CNS, it might well be related to a reduction in viscosity of the blood thereby improving memory, as for example in case of *Ginkgo biloba* L. (Santos et al., 2003).

Plants not identified up to the species level in the books were not included in the study. This decision was based on the fact that the attempt to establish a correlation between the plants cited only by their vernacular names or by genus and possible

Table 1  
Thirty-four species native to Brazil cited in historic literature with possible uses/effects on the central nervous system

Species (family)	Vernacular name	Indications in historic literature [part utilized]	Indications by Brazilian communities nowadays
<i>Anacardium occidentale</i> L. (Anacardiaceae)	Cajú (cashew)	“Intoxicating, inebriating” [fruit] (Martius, 1939)	Hemorrhoids, serious diarrhea [bark]—coastal Caiçara fishermen (Di Stasi and Hiruma-Lima, 2002); lower extremity pain, skin injury [bark]—Pataxó Indians (Thomas, 2001); to treat inflammations [bark] (Albuquerque and Andrade, 2002); inflammation—Xucuru Indians (Silva and Andrade, 1998)
<i>Aristolochia grandiflora</i> Sw (Aristolochiaceae)	Mil-homens	“Febrifuge” [stem] (Gomes, 1972)	No data
<i>Bertholletia excelsa</i> Bompl. (Lecythidaceae)	Castanhero-do-pará (brazil-nut-tree)	“Smoke” [bark] (Spix and Martius, 1976)	Anemia [fruit]—Amazonian river-dwellers (Amorozo and Gély, 1988); gastrointestinal disturbances, fever, body strengtheners [bark and fruit]—Amazonian river-dwellers (Rodrigues, 2006); to treat dysentery, colic, liver [bark and fruit]—Amazonian rubber tappers (Ming, 1995)
<i>Bignonia chica</i> Humb. and Bonpl. (Bignoniaceae)	Carajuru	“Blood purifier” [leaf] (Spix and Martius, 1976)	Gastrointestinal disturbances, body strengtheners [leaf]—Amazonian river-dwellers (Rodrigues, 2006); conjunctivitis [leaf]—Amazonian inhabitants (Estrella, 1995)
<i>Cecropia palmata</i> Willd. (Cecropiaceae)	No record	“Becalmes the need for food and sleep” [leaf] (Spix and Martius, 1976)	No data
<i>Couratari guianensis</i> Aubl. (Lecythidaceae)	Tauiri	“Cigarette” [bark] (Spix and Martius, 1976)	No data
<i>Cryptocarya pretiosa</i> Mart. ex Nees (Lauraceae)	Casca-preciosa	“To counteract weakness in the nerves” [bark] (Spix and Martius, 1976)	No data
<i>Dorstenia brasiliensis</i> Lam. (Moraceae) <sup>a</sup>	Contraheerva	“Febrifuge” [root and leaf] (Gomes, 1972)	To treat snake bites, fever and others infections (Lorenzi and Matos, 2002)
<i>Echites cururu</i> Mart. (Apocynaceae)	Cipó-cururu	“Counteracts fever, headache, and loss of appetite” [stem] (Spix and Martius, 1976; Martius, 1939)	No data
<i>Echites grandiflorus</i> G. Mey. (Apocynaceae)	Cipó-cururu	“Counteracts fever, headache, and loss of appetite” [stem] (Spix and Martius, 1976)	No data
<i>Erythroxylon coca</i> Lam. (Erythroxylaceae) <sup>a</sup>	Ipadú	“Inebriates” [leaf] (Spix and Martius, 1976; Martius, 1939)	Stimulant and narcotic—Amazonian Indians (Plowman, 1979; Cooper, 1987), to treat heart disorders—Amazonian Inhabitants (Van den Berg and Silva, 1988)
<i>Hura crepitans</i> L. (Euphorbiaceae)	Assacú	“Cigarette (numbs)” [sap] (Spix and Martius, 1976)	Ray sting [latex]—Amazonian river-dwellers (Amorozo, 1993); ichthyotoxic (Heizer, 1987; Schultes and Raffaulf, 1990)
<i>Manihot utilisima</i> Pohl (Euphorbiaceae)	Mandioca (manioc)	“Intoxicates, inebriating” [root] (Martius, 1939)	No data
<i>Musa sapientum</i> L. (Musaceae)	Banana	“Intoxicates, inebriating” [fruit] (Martius, 1939)	No data
<i>Nicotiana langsdorffii</i> Weinmann (Solanaceae)	Nicotina (nicotine)	“Same use as <i>Nicotiana tabacum</i> and <i>N. rustica</i> (narcotic)” [whole plant] (Martius, 1939)	No data
<i>Ocotea kujumari</i> Mart. (Lauraceae)	Cujumari	“Combats rheumatic pain” [seed] (Spix and Martius, 1976)	No data
<i>Ocotea opifera</i> Mart. (Lauraceae)	No record	“Combats rheumatic pain” [seed] (Spix and Martius, 1976)	“Enhances the ‘strength’ of the coca” [fruit]—Amazonian Indians (Schultes and Raffaulf, 1990)
<i>Paullinia cururu</i> L. (Sapindaceae)	Guaraná	“Affects the emotions, uncommon excitement, double vision, insomnia, aphrodisiacs” [seed] (Spix and Martius, 1976)	No data

Table 1 (Continued)

Species (family)	Vernacular name	Indications in historic literature [part utilized]	Indications by Brazilian communities nowadays
<i>Paullinia pinnata</i> L. (Sapindaceae)	Guaraná/Curuapé/ Cruapé-Vermelho	“Affects emotions, uncommon excitement, double vision, insomnia, aphrodisiac” [seed] (Spix and Martius, 1976)	No data
<i>Paullinia sorbilis</i> Mart. (Sapindaceae)	Guaraná	“Affects the emotions, uncommon excitement, double vision, insomnia, aphrodisiac” [seed] (Spix and Martius, 1976)	No data
<i>Piper nodosum</i> C. DC. (Piperaceae)	No record	“Combats toothache” [root] (Martius, 1939)	No data
<i>Piptadenia colubrina</i> (Vell.) Benth. (Fabaceae)	Paricá	“Tonic” [seed] (Martius, 1939)	No data
<i>Psychotria ipecacuanha</i> (Brot.) Stokes (Rubiaceae)	Ipecacuanha	“Antidote to opium” [root] (Gomes, 1972)	Antiemetic [root] (Lorenzi and Matos, 2002); stimulant (snuff)—Amazonian Indians (Cooper, 1987)
<i>Selaginella convoluta</i> (Arn.) Spring (Selaginellaceae)	No record	“Arouses vital forces that are dormant (analeptic)” [whole plant] (Martius, 1939)	No data
<i>Tachia guianensis</i> Aubl. (Gentianaceae)	Raiz-de-jacaré- aru/coferana	“Febrifuge” [root] (Spix and Martius, 1976)	To treat “sore stomach” [root]—Amazonian Indians (Schultes and Raffauf, 1990)
<i>Theobroma bicolor</i> Bonpl. (Sterculiaceae)	Cacau (cacao)	“Bitter principle comparable to caffeine” [whole plant] (Spix and Martius, 1976)	No data
<i>Theobroma cacao</i> L. (Sterculiaceae)	Cacau (cacao)	“Bitter principle comparable to caffeine” [whole plant] (Spix and Martius, 1976)	Osteomuscular problems [fruit]—Amazonian river-dwellers (Rodrigues, 2006); to treat liver problems—Amazon Inhabitants (Van den Berg and Silva, 1988)
<i>Theobroma microcarpum</i> Mart. (Sterculiaceae)	Cacau (cacao)	“Bitter principle comparable to caffeine” [whole plant] (Spix and Martius, 1976)	No data
<i>Theobroma speciosum</i> Willd. ex Spreng. (Sterculiaceae) <sup>a</sup>	Cacau (cacao)	“Bitter principle comparable to caffeine” [whole plant] (Spix and Martius, 1976)	Diuretic, stimulant [seed] (Lorenzi and Matos, 2002)
<i>Theobroma subincanum</i> Martius in Buchner (Sterculiaceae)	Cacau (cacao)	“Bitter principle comparable to caffeine” [whole plant] (Spix and Martius, 1976)	Hallucinogenic snuff [bark ashes]—Indians living in the Northwestern Amazon (Schultes, 1969)
<i>Theobroma sylvestre</i> Aubl. ex Mart. in Buchner (Sterculiaceae)	Cacau (cacao)	“Bitter principle comparable to caffeine” [whole plant] (Spix and Martius, 1976)	No data
<i>Trimezia lurida</i> Salisb. (Iridaceae)	Manciçó	“Febrifuge” [root] (Gomes, 1972)	No data
<i>Vandellia difusa</i> L. (Scrophulariaceae)	Erva-da-mata-cana	“Febrifuge” [root] (Spix and Martius, 1976)	No data
<i>Zea mays</i> L. (Poaceae)	Milho (maize)	“Inebriates” [fruit] (Martius, 1939)	Antiemetic [straw-fibers]—Terena Indians (Carvalho, 1996); kidney pain, measles [straw-fibers]—Amazonian rubber tappers (Ming, 1995)

<sup>a</sup> Species that present coincidence between the uses described in historic literature and in the current ethnopharmacological studies.

species would be of little value since the descriptive material on these species did not contain enough information to distinguish precisely to what species the authors were referring. Thus, of the 93 plants recorded, 39 were excluded from the present study, because they were described by their vernacular names or were identified on a genus level, while 54 were identified on a species level. Of these 54 plants, 34 are native to Brazil and the results referred in the present article are related to these species (Tables 1 and 2).

Ethnopharmacological studies showed that only 13 of the 34 species have been recorded in modern ethnopharmacological studies in traditional communities (Table 1). Most of the uses reported in former centuries do not coincide with those today.

Only three cases were reported in which a use coincided as can be noted (<sup>a</sup>) in Table 1. Thus, *Dorstenia brasiliensis* Lam. and *Theobroma speciosum* Willd. ex Spreng. were cited for their febrifuge and stimulating action, respectively (Lorenzi and Matos, 2002) and these uses were also reported in previous centuries. Also, *Erythroxylon coca* Lam. has been cited as “inebriating” in the nineteenth century (Martius, 1939; Spix and Martius, 1976), and nowadays its stimulant and narcotic effects have been reported by Plowman (1979) and Cooper (1987), among Amazonian Indians.

A bibliographical survey showed that up to the present time, only eight of these 34 species were the focus of pharmacological investigation (Table 2). Two coincidences between uses reported

Table 2  
Thirty-four species native to Brazil cited in historic literature with possible uses/effects on the central nervous system, as well as their phytochemical and pharmacological studies found in present-day literature

Species (family)	Phytochemical data found in present-day literature	Pharmacological studies found in present-day literature	Pharmacological use in the patent
<i>Anacardium occidentale</i> L. (Anacardiaceae)	Anacardic acids, cardanols and cardols (Trevisan et al., 2006; Olatunji et al., 2005); steroids (Phillips et al., 2005; Alexander-Lindo et al., 2004); terpenes, tannins, flavonoids and saponins (Gonçalves et al., 2005); carotenoids and ascorbic acid (Assunção and Mercadante, 2003); ketones, lactones and norisoprenoids (Bicalho et al., 2000); tannins (Mota et al., 1985); long-chain phenols (Tyman et al., 1984); acylated flavanone glycoside (Rahman et al., 1978)	Antioxidant (Melo Cavalcante et al., 2003; Trevisan et al., 2006); anti-inflammatory (Mota et al., 1985); anti-diabetic effects (Olatunji et al., 2005)	Antioxidant, anti-obesity
<i>Aristolochia grandiflora</i> Sw (Aristolochiaceae)	No data	No data	No data
<i>Bertholletia excelsa</i> Bompl. (Lecythidaceae)	Saponin, phenolic compounds and quinazolinic alkaloids (Campos et al., 2005); amyrynes and tocopherols (De Siqueira et al., 2003)	No data	No data
<i>Bignonia chica</i> Humb. and Bonpl. (Bignoniaceae)	No data	No data	No data
<i>Cecropia palmata</i> Willd. (Cecropiaceae)	No data	No data	No data
<i>Couratari guianensis</i> Aubl. (Lecythidaceae)	No data	No data	No data
<i>Cryptocarya pretiosa</i> Mart. ex Nees (Lauraceae)	No data	No data	No data
<i>Dorstenia brasiliensis</i> Lam. (Moraceae)	Seco-adianane-type triterpenoids, isopimarane-type diterpenoid and coumarins (Uchiyama et al., 2002); steroids and furocoumarins (Kuster et al., 1994)	No data	No data
<i>Echites cururu</i> Mart. (Apocynaceae)	No data	No data	No data
<i>Echites grandiflorus</i> G. Mey. (Apocynaceae)	No data	No data	No data
<i>Erythroxylon coca</i> Lam. (Erythroxylaceae)	Tropane alkaloids (Hammerschmidt, 1999; Leete, 1983); alkaloids (Turner et al., 1981)	Anesthetic (Bedford et al., 1984)	Anorectic
<i>Hura crepitans</i> L. (Euphorbiaceae)	Flavonoids (Freixa et al., 1998)	Ichthyotoxic (Pires, 1978)	Neuronal regeneration
<i>Manihot utilisima</i> Pohl (Euphorbiaceae)	Flavonoids (Chariandy et al., 1999; Kamil et al., 1974)	Antioxidant (Rahmat et al., 2004)	No data
<i>Musa sapientum</i> L. (Musaceae)	Essential oils (Pino et al., 2003; Pannangpetch et al., 2001); flavonoids (Lewis et al., 1999; Pari and Maheswari, 1999); cycloartane triterpenes (Akihisa et al., 1998); tannins (Ali and Bhutani, 1993); steroids (Knapp et al., 1972)	Antioxidant (Goel et al., 2001); muscle paralysis (Singh and Dryden, 1985)	Antidepressant
<i>Nicotiana langsdorffii</i> Weinmann (Solanaceae)	No data	No data	No data
<i>Ocotea cujumary</i> Mart. (Lauraceae)	No data	No data	No data
<i>Ocotea opifera</i> Mart. (Lauraceae)	Essential oils (Lorenzo et al., 2001)	No data	No data
<i>Paullinia cururu</i> L. (Sapindaceae)	No data	No data	No data
<i>Paullinia pinnata</i> L. (Sapindaceae)	Polyphenols (Zamble et al., 2006); flavone glycosides (Abourashed et al., 1999)	Ichthyotoxic (Pires, 1978)	No data
<i>Paullinia sorbilis</i> Mart. (Sapindaceae)	No data	No data	Antidepressant
<i>Piper nodosum</i> C. DC. (Piperaceae)	No data	No data	No data



Table 2 (Continued)

Species (family)	Phytochemical data found in present-day literature	Pharmacological studies found in present-day literature	Pharmacological use in the patent
<i>Piptadenia colubrina</i> (Vell.) Benth. (Fabaceae)	Indole alkaloids (Pachter et al., 1959)	No data	No data
<i>Psychotria ipecacuanha</i> (Brot.) Stokes (Rubiaceae)	Alkaloids (Garcia et al., 2005)	No data	No data
<i>Selaginella convoluta</i> (Am.) Spring (Selaginellaceae) <sup>a</sup>	No data	Antidepressant (Santos et al., 1994)	No data
<i>Tachia guianensis</i> Aubl. (Gentianaceae)	Flavonoids (Carvalho et al., 1991)	No data	No data
<i>Theobroma bicolor</i> Bonpl. (Sterculiaceae)	Fatty acids (Jee, 1984); phenolic acids (Torres et al., 2002)	No data	No data
<i>Theobroma cacao</i> L. (Sterculiaceae) <sup>a</sup>	Catechins (Gotti et al., 2006; Kelm et al., 2006); flavonoids (Ramiro et al., 2005; Osman et al., 2004); flavonoids O-glucosides and C-glucosides (Sanchez-Rabeneda et al., 2003); tannins (Falade et al., 2005); essential oils (Chee et al., 2005); alkaloids (Stark et al., 2005); polyphenols (Edwards et al., 2005); proanthocyanidin glycosides (Hatano et al., 2002); carboxylic acids, purine alkaloids, fatty acids (Bucheli et al., 2001)	Anti-inflammatory (Schewe et al., 2002); psychoactive (Melzig et al., 2000)	Anti-inflammatory; antioxidant; appetite regulator
<i>Theobroma microcarpum</i> Mart. (Sterculiaceae)	No data	No data	No data
<i>Theobroma speciosum</i> Willd. ex Spreng. (Sterculiaceae)	No data	No data	No data
<i>Theobroma subincanum</i> Martius in Buchner (Sterculiaceae)	Tocopherol, fatty acids, steroids (Bruni et al., 2000, 2002)	No data	No data
<i>Theobroma sylvestre</i> Aubl. ex Mart. in Buchner (Sterculiaceae)	No data	No data	No data
<i>Trimezia lurida</i> Salisb. (Iridaceae)	No data	No data	No data
<i>Vandellia difusa</i> L. (Scrophulariaceae)	No data	No data	No data
<i>Zea mays</i> L. (Poaceae)	Terpenoids (Yan and Wang, 2006); polyamines (Moreau and Hicks, 2005); fatty acids, primary alcohols, diacids, omega-hydroxy fatty acids, and 2-hydroxy fatty acids, with omega-hydroxy fatty acids (Schreiber et al., 2005); phenolic acids (Fontaine et al., 2005; Funk et al., 2005); alkylresorcinol (Gembeh et al., 2001); sesquiterpene hydrocarbon (Köllner et al., 2004); polyphenols (Del Pozo-Insfran et al., 2006)	No data	No data

<sup>a</sup> Species that present coincidence between the uses described in historic literature and their effects observed in the nowadays pharmacological studies.

in historic literature and the pharmacological data described in contemporary scientific literature were noted also in Table 2. *Selaginella convoluta* (Arn.) Spring and *Theobroma cacao* L. were cited in historic literature for its arousing “vital forces that are dormant (analeptic)” (Martius, 1939) and for its “Bitter principle comparable to caffeine” (Spix and Martius, 1976), respectively (Table 1). The current pharmacological studies showed antidepressant (Santos et al., 1994) and psychoactive (Melzig et al., 2000) effects, respectively (Table 2).

Of the 34 species cited in the present study, only 16 were studied from a phytochemical point of view (Table 2). The

main chemical constituents found in these plants in decreasing order were: flavonoids (eight plants), alkaloids (six), phenolic acids (four), tannins (four) essential oils (three), pentacyclic triterpenes (three), steroids (two) and saponins (two).

In the survey on possible requests for patents, six species were found to be registered (*Anacardium occidentale* L., *Erythroxylon coca* Lam., *Hura crepitans* L., *Musa sapientum* L., *Paullinia sorbilis* Mart., and *Theobroma cacao* L.). However, no patent shows any coincidence in plant use and the possible effects described in historic literature (Tables 1 and 2).

No information was found concerning medication registered in Brazil and developed as from these plants.

#### 4. Discussion

This study presents data that allows us to verify not only ethnopharmacological data but also social and cultural aspects of Brazilian cultures of the past century, in compliance with a recent ethnopharmacological concept as a transdisciplinary exploration that spans the biological and social sciences (Etkin and Elisabetsky, 2005).

Some limitations must be considered in spite of the careful survey. For instance, use of a scientific name as a key word in databases will not bring results if the orthography of the scientific name published has been subject to alteration. Also, it was not checked on all the botanic synonyms in this survey. Finally, maybe not all Brazilian studies have been consulted as desired, since most of them have been published in periodic not indexed.

From the ethnopharmacological point of view, the results showed the different paths traveled by plants over the course of history, confirming the hypothesis that some of them were used for the same purposes for centuries (*Dorstenia brasiliensis* Lam.; *Erythroxylon coca* Lam. and *Theobroma speciosum* Willd. ex Spreng.). For others, uses were modified, as can be observed in the recent ethnopharmacological studies described in Table 1 (*Anacardium occidentale* L.; *Bertholletia excelsa* Bompl.; *Bignonia chica* Humb. and Bonpl.; *Hura crepitans* L.; *Ocotea opifera* Mart.; *Psychotria ipecacuanha* (Brot.) Stokes; *Tachia guianensis* Aubl.; *Theobroma cacao* L.; *Theobroma subincanum* Martius in Buchner and *Zea mays* L.). In other cases, uses were completely forgotten in the course of time (*Aristolochia grandiflora* Sw; *Cecropia palmata* Willd.; *Couratari guianensis* Aubl.; *Cryptocarya pretiosa* Mart., ex Nees; *Echites cururu* Mart.; *Echites grandiflorus* G. Mey.; *Manihot utilissima* Pohl; *Musa sapientum* L.; *Nicotiana langsdorffii* Weinmann; *Ocotea cujumary* Mart.; *Paullinia cururu* L.; *Paullinia pinnata* L.; *Paullinia sorbilis* Mart.; *Piper nodosum* C. DC.; *Piptadenia colubrina* (Vell.) Benth.; *Selaginella convoluta* (Arn.) Spring; *Theobroma bicolor* Bonpl.; *Theobroma microcarpum* Mart.; *Theobroma seylvestre* Aubl. ex Mart. in Buchner; *Trimezia lurida* Salisb. and *Vandellia difusa* L.).

From the phytochemical point of view, the most recurrent chemical constituents in the plants under study were the flavonoids, alkaloids, phenolic acids, tannins and essential oils. As has been reported in several articles, these substances exert activity on the CNS. Some flavonoids present anticonvulsive and sedative effects (Dos Santos et al., 2005; Fernandez et al., 2004), have the effect of protecting against neurodegeneration of the CNS (Nagase et al., 2005; Heitzman et al., 2005; Marder et al., 2003) and antidepressant activity (Juergenliemk et al., 2003). Many tannins showed an anticonvulsive effect (Dos Santos et al., 2005). Pentacyclic triterpenes also presented activity on the CNS (Heitzman et al., 2005). Alkaloids of the matrine type exert an antinociceptive effect through many mechanisms of action (Yin and Zhu, 2005), such as an increase in activation of the cholinergic system (Breining, 2004). Alkaloids thus possess pharmacological activity and may be relevant in the treatment

of cognitive disorders, including an increase in the cholinergic function in the CNS (Howes and Houghton, 2003).

Phenolic compounds protect neurons from oxidative stress (Scapagnini et al., 2004). Aromatic plants possess high concentrations of essential oils and are utilized in the treatment of diseases related to the CNS, common in tropical regions (Freire et al., 2006; Ozturk et al., 2002).

The reduced number of pharmacological studies found for the species surveyed shows that they obviously did not focus on medicinal plants cited in historic literature. Hopefully this article contributes to a new appreciation of such historical documents, since researches which reinstate historic medicinal uses of plants are rare or difficult to carry through (because of difficult access to books), so that few species have been the subject of investigation up to the present time. The main difficulty concerns to the lack of a location which concentrates these books or even information about them, since they are dispersed in few museums and other institutions; often in bad conditions of conservation. Also, sometimes they are only available as microfilms, and usually are contaminated by fungus, jeopardizing the reading. A wider search has been conducted by the authors in other Brazilian cities, in order to verify the general conditions of the historic literature deposited in Institutions outside Sao Paulo.

Only six species were found to have a registered patent in the patent database consulted (Table 2). With the exception of *Paullinia sorbilis* Mart., all five other plants have been studied from a pharmacological point of view. *Paullinia sorbilis* Mart. was also the only species related to the CNS with a patent registered in Brazil (INPI, 2005); all others have been patented in the United States or in Europe. In Brazil, only methods or inventions are patentable—not species and their constituents. When a substance is isolated, or effects of one particular species reported, this register can only be made outside Brazil, in countries that accept this type of patent. Many species are taken to other countries and studied in foreign laboratories. These two facts may explain the number of patents of Brazilian species registered in international offices and the absence of patents for Brazilian species in the country.

The fact that no records have been found of medications in Brazil for the species under study comes as no surprise. With the exception of one phytomedicine registered by ANVISA in 2004, Acheflan<sup>®</sup>, Brazil does not possess any registered phytomedicine derived from Brazilian resources. This is due to several reasons, ranging from the lack of financial incentive from the government for the study of natural products, to the recent difficulties in the Brazilian legislation concerning the study of medicinal plants. Some of these obstacles have been reverted over the last years. Firstly, through incentive provided by university-business company partnerships, financing wide-scale projects and training groups of professionals of great importance to Brazil (Rodrigues, 2005). Secondly, registration of one particular phytomedicine takes years to pass all of the pre-clinical and clinical pharmacological tests. For such, the phytomedicine would have to show its pharmacological safety and effectiveness – in itself, not a very simple matter – since many studies are not carried through because the species do not

present the desired pharmacological effect in the first pharmacological tests.

## 5. Conclusion

This study shows the importance and ethnopharmacological potential of bibliographical surveys of historic texts in providing a record of plants with possible bioactive properties, often forgotten over the course of time. The species of greatest importance are those which have not been the object of pharmacological investigation in spite of their historic uses reported.

## Acknowledgements

Our grateful acknowledgements to FAPESP for financial support, to Dr. E.A. Carlini for all his support and guidance, to Dr. Amélia Vera Guimarães de Sousa for her support, and to Dr. Zuleika Ribeiro do Valle, for kindly lending us the books used in this study.

## References

- Abourashed, E.A., Toyang, N.J., Choinski, J., Khan, I.A., 1999. Two new flavone glycosides from *Paullinia pinnata*. *Journal of Natural Products* 62, 1179–1181.
- Akihisa, T., Kimura, Y., Tamura, T., 1998. Cycloartane triterpenes from the fruit peel of *Musa sapientum*. *Phytochemistry* 47, 1107–1110.
- Albuquerque, U.P., Andrade, L.H.C., 2002. Uso de Recursos Vegetais da Caatinga: o caso do Agreste do Estado de Pernambuco (Nordeste do Brasil). *Interciencia* 27, 336–346.
- Alexander-Lindo, R.L., Morrison, E.Y.S., Nair, M.G., 2004. Hypoglycaemic effect of stigmast-4-en-3-one and its corresponding alcohol from the bark of *Anacardium occidentale* (Cashew). *Phytotherapy Research* 18, 403–407.
- Ali, M., Bhutani, K.K., 1993. Flavan-3,4-diols from *Musa sapientum* seeds. *Pharmazie* 48, 455–456.
- Amorozo, M.C.M., Gély, A., 1988. Uso de Plantas Mediciniais por Caboclos do Baixo Amazonas. *Boletim do Museu paraense Emilio Goeldi*, vol. 4. Bacarena, PA, Brasil, pp. 47–135.
- Amorozo, M.C.M., 1993. Algumas Notas Adicionais Sobre o Emprego de Plantas e Outros Produtos com fins terapeuticos Pela População Cabocla do Município de Bacarena, PA, Brasil. *Boletim do Museu paraense Emilio Goeldi* 9, 249–268.
- ANVISA—Agenda Nacional de Vigilância Sanitária, 2005. <http://www.anvisa.gov.br/> (accessed 24–26, July 2005).
- Assunção, R.B., Mercadante, A.Z., 2003. Carotenoids and ascorbic acid composition from commercial products of cashew apple (*Anacardium occidentale* L.). *Journal of Food Composition and Analysis* 16, 647–657.
- Bedford, J.A., Turner, C.E., Elsohly, H.N., 1984. Local anesthetic effects of cocaine and several extracts of the coca leaf (*E. coca*). *Pharmacology Biochemistry and Behavior* 20, 819–821.
- Bicalho, B., Pereira, A.S., Neto, F.R.A., Pinto, A.C., Rezende, C.M., 2000. Application of high-temperature gas chromatography–mass spectrometry to the investigation of glycosidically bound components related to cashew apple (*Anacardium occidentale* L. Var. nanum) volatiles. *Journal of Agricultural and Food Chemistry* 48, 1167–1174.
- Breining, S.R., 2004. Recent developments in the synthesis of nicotinic acetylcholine receptor ligands. *Current Topics in Medicinal Chemistry* 4, 609–629.
- Bruni, R., Bianchini, E., Bettarello, L., Sacchetti, G., 2000. Lipid composition of wild Ecuadorian *Theobroma subincanum* Mart. seeds and comparison with two varieties of *Theobroma cacao* L. *Journal of Agricultural and Food Chemistry* 48, 691–694.
- Bruni, R., Medici, A., Guerrini, A., Scalia, S., Poli, F., Romagnoli, C., Muzzoli, M., Sacchetti, G., 2002. Tocopherol, fatty acids and sterol distributions in wild Ecuadorian *Theobroma subincanum* (Sterculiaceae) seeds. *Food Chemistry* 77, 337–341.
- Bucheli, P., Rousseau, G., Alvarez, M., Laloi, M., McCarthy, J., 2001. Developmental variation of sugars, carboxylic acids, purine alkaloids, fatty acids, and endoproteinase activity during maturation of *Theobroma cacao* L. seeds. *Journal of Agricultural and Food Chemistry* 49, 5046–5051.
- Carvalho, L.H., Brandão, M.G.L., Santos, D., Lopes, J.L.C., Krettli, A.U., 1991. Antimalarial activity of crude extracts from Brazilian plants studied in vivo in *Plasmodium berghei*-infected mice and in vitro against *Plasmodium falciparum* in culture. *Brazilian Journal of Medical and Biological Research* 24, 1113–1123.
- Campos, F.R., Januario, A.H., Rosas, L.V., Nascimento, S.K.R., Pereira, P.S., Franca, S.C., Cordeiro, M.S.C., Toldo, M.P.A., Albuquerque, S., 2005. Trypanocidal activity of extracts and fractions of *Bertholletia excelsa*. *Fitoterapia* 76, 26–29.
- Carvalho, F., 1996. Koixomuneti e outros curadores: xamanismo e práticas de cura entre os Terena. Doctorate Thesis. University of Sao Paulo, Sao Paulo, 146 p.
- Chariandy, C.M., Seaforth, C.E., Phelps, R.H., Pollard, G.V., Khambay, B.P.S., 1999. Screening of medicinal plants from Trinidad and Tobago for antimicrobial and insecticidal properties. *Journal of Ethnopharmacology* 64, 265–270.
- Chee, S.Y.K., Malek, S.N.A., Ramli, N., 2005. Essential oils in the leaves of cocoa (*Theobroma cacao* L.) clone UIT1 and NA33. *Journal of Essential Oil Research* 17, 312–313.
- Cooper, J.M., 1987. Estimulantes e Narcóticos. In: Ribeiro, D., et al. (Eds.), *Handbook of South American Indians*. FINEP, Petrópolis.
- DEF (Dicionário de Especialidades Farmacêuticas), 2004/2005. *Jornal Brasileiro de Medicina*, Sao Paulo.
- De Siqueira, D.S., Pereira, A.D., Neto, F.R.D., Cabral, J.A., Ferreira, C.A.C., Simoneit, B.R.T., Elias, V.O., 2003. Determination of high molecular mass compounds from Amazonian plant's leaves. *Quimica Nova* 26, 633–640.
- Del Pozo-Insfran, D., Brenes, C., Saldívar, S.O.S., Talcott, S.T., 2006. Polyphenolic and antioxidant content of white and blue corn (*Zea mays* L.) products. *Food Research International* 39, 696–703.
- Di Stasi, L.C., Hiruma-Lima, C.A., 2002. Plantas Mediciniais na Amazônia e na Mata Atlântica. UNESP, Sao Paulo.
- Dos Santos, J.G., Blanco, M.M., Do Monte, F.H.M., Russi, M., Lanzotti, V.M.N.B., Leal, L.K.A.M., Cunha, G.M., 2005. Sedative and anticonvulsant effects of hydroalcoholic extract of *Equisetum arvense*. *Fitoterapia* 76, 508–513.
- Duke, 2006. Dr. Duke's Phitochemical and Ethnobotanical Databases, <http://www.ars-grin.gov/duke/> (accessed May, 2006).
- Edwards, H.G.M., Villar, S.E.J., de Oliveira, L.F.C., Le Hyaric, M., 2005. Analytical Raman spectroscopic study of cacao seeds and their chemical extracts. *Analytica Chimica Acta* 538, 175–180.
- ESPACENET (European Patente Office), 2005. <http://www.ep.espacenet.com/> (accessed August 15, 2005).
- Estrella, E., 1995. *Plantas Medicinales Amazonicas: Realidad y Perspectivas*. Tratado de Cooperación Amazonica. TCA, Lima.
- Etkin, N.L., Elisabetsky, E., 2005. Seeking a transdisciplinary and culturally germane science: the future of ethnopharmacology. *Journal of Ethnopharmacology* 100, 23–26.
- Fabre, A.J., 2003. Mythology and the medicinal plants of antiquity. *Histoire des Sciences Medicales* 37, 65–87.
- Falade, O.S., Otemuyiwa, I.O., Oladipo, A., Oyedapo, O.O., Akinpelu, B.A., Adewusi, S.R.A., 2005. The chemical composition and membrane stability activity of some herbs used in local therapy for anemia. *Journal of Ethnopharmacology* 102, 15–22.
- Fernandez, S., Wasowski, C., Paladini, A.C., Marder, M., 2004. Sedative and sleep-enhancing properties of linarin, a flavonoid-isolated from *Valeriana officinalis*. *Pharmacology Biochemistry and Behavior* 77, 399–404.
- Fontaine, A.S., Bout, S., Barriere, Y., Vermerris, W., 2005. Variation in cell wall composition among forage maize (*Zea mays* L.) inbred lines and its impact on digestibility: analysis of neutral detergent fiber composition by pyrolysis–gas chromatography–mass spectrometry. *Journal of Agricultural and Food Chemistry* 51, 8080–8087.



- Freire, C.M.M., Marques, M.O.M., Costa, M., 2006. Effects of seasonal variation on the central nervous system activity of *Ocimum gratissimum* L. essential oil. *Journal of Ethnopharmacology* 105, 161–166.
- Freixa, B., Vila, R., Vargas, L., Lozano, N., Adzet, T., Canigual, S., 1998. Screening for antifungal activity of nineteen Latin American plants. *Phytotherapy Research* 12, 427–430.
- Funk, C., Ralph, J., Steinhart, H., Bunzel, M., 2005. Isolation and structural characterisation of 8-O-4/8-O-4- and 8-8/8-O-4-coupled dehydrotriferulic acids from maize bran. *Phytochemistry* 66, 363–371.
- García, R.M.A., de Oliveira, L.O., Moreira, M.A., Barros, W.S., 2005. Variation in emetine and cephaeline contents in roots of wild Ipecac (*Psychotria ipecacuanha*). *Biochemical Systematics and Ecology* 33, 233–243.
- Gembah, S.V., Brown, R.L., Grimm, C., Cleveland, T.E., 2001. Identification of chemical components of corn kernel pericarp wax associated with resistance to *Aspergillus flavus* infection and aflatoxin production. *Journal of Agricultural and Food Chemistry* 49, 4635–4641.
- Goel, R.K., Sairam, K., Rao, C.V., 2001. Role of gastric antioxidant and anti-*Helicobacter pylori* activities in antiulcerogenic activity of plantain banana (*Musa sapientum* var. *paradisica*). *Indian Journal of Experimental Biology* 39, 719–722.
- Gomes, B.A., 1972. *Plantas Medicinais do Brasil (1801–1812)*, vol. 5. *Brasilien-sia Documenta*, Sao Paulo.
- Gonçalves, J.L.S., Lopes, R.C., Oliveira, D.B., Costa, S.S., Miranda, M.M.F.S., Romanos, M.T.V., Santos, N.S.O., Wigg, M.D., 2005. In vitro anti-rotavirus activity of some medicinal plants used in Brazil against diarrhea. *Journal of Ethnopharmacology* 99, 403–407.
- Gotti, R., Furlanetto, S., Pinzauti, S., Cavrini, V., 2006. Analysis of catechins in *Theobroma cacao* beans by cyclodextrin-modified micellar electrokinetic chromatography. *Journal of Chromatography A* 1112, 345–352.
- Hammerschmidt, D.E., 1999. *Erythroxylon coca*: cocaine and norcocaine. *The Journal of Laboratory and Clinical Medicine* 133, 613–614.
- Hatano, T., Miyatake, H., Natsume, M., Osakabe, N., Takizawa, T., Ito, H., Yoshida, T., 2002. Proanthocyanidin glycosides and related polyphenols from cacao liquor and their antioxidant effects. *Phytochemistry* 59, 749–758.
- Heinrich, M., Teoh, H.L., 2004. Galanthamine from snowdrop—the development of a modern drug against Alzheimer's disease from local Caucasian knowledge. *Journal of Ethnopharmacology* 92, 147–162.
- Heinrich, M., Kufer, J., Leonti, M., Pardo-de-Santayana, M., 2006. Ethnobotany and Ethnopharmacology—Interdisciplinary links with the historical sciences. *Journal of Ethnopharmacology* 107, 157–160.
- Heitzman, M.E., Neto, C.C., Winiarz, E., Vaisberg, A.J., Hammond, G.B., 2005. Ethnobotany, phytochemistry and pharmacology of *Uncaria* (Rubiaceae). *Phytochemistry* 66, 5–29.
- Heizer, R.F., 1987. Venenos de pesca. In: Ribeiro, D., et al. (Eds.), *Handbook of South American Indians*. FINEP, Petrópolis, pp. 159–171.
- Howes, M.J.R., Houghton, P.J., 2003. Plants used in Chinese and Indian traditional medicine for improvement of memory and cognitive function. *Pharmacology Biochemistry and Behavior* 75, 513–527.
- INPI—Instituto Nacional de Propriedade Industrial, 2005. <http://www.inpi.gov.br/> (accessed 13, August 2005), International Society of Ethnopharmacology, 2005. ISE Constitution. <http://www.ethnopharmacology.org/iseConst.html> (accessed 31, May 2006).
- Jee, M.H., 1984. Composition of the fat extracted from the seeds of *Theobroma bicolor*. *Journal of the American Oil Chemists' Society* 61, 751–753.
- Juergeniemk, G., Boje, K., Huewel, S., Lohmann, C., Galla, H.J., Nahrstedt, A., 2003. In vitro studies indicate that miquelianin (quercetin 3-O-beta-D-glucuronopyranoside) is able to reach the CNS from the small intestine. *Planta Medica* 69, 1013–1017.
- Kamil, M., Ilyas, M., Rahman, W., Okigawa, M., Kawano, N., 1974. Biflavones from *Manihot utilisima*. *Phytochemistry* 13, 2619–2620.
- Kelm, M.A., Johnson, J.C., Robbins, R.J., Hammerstone, J.F., Schmitz, H.H., 2006. High-performance liquid chromatography separation and purification of cacao (*Theobroma cacao* L.) procyanidins according to degree of polymerization using a diol stationary phase. *Journal of Agricultural and Food Chemistry* 54, 1571–1576.
- Knapp, F.F., Phillips, D.O., Goad, L.J., Goodwin, T.W., 1972. Isolation of 14alpha-methyl-9beta,19-cyclo-5alpha-ergost-24(28)-en-3beta-ol from *Musa sapientum*. *Phytochemistry* 2, 3497–3500.
- Köllner, T.G., Schnee, C., Gershenzon, J., Degenhardt, J., 2004. The sesquiterpene hydrocarbons of maize (*Zea mays*) form five groups with distinct developmental and organ-specific distributions. *Phytochemistry* 65, 1895–1902.
- Kufer, J., Förther, H., Pöll, E., Heinrich, M., 2005. Historical and modern medicinal plant uses—the example of the Chiortii Maya and Ladinos in Eastern Guatemala. *Journal of Pharmacy and pharmacology* 57, 1127–1152.
- Kury, L., 2001. Viajantes-naturalistas no Brasil oitocentista: experiência, relato e imagem. *História, Ciência, Saúde-Manguinhos* vol. 8 (Suppl.), 863–880.
- Kuster, R.H., Bernardo, R.R., Da Silva, A.J.R., Parente, J.P., Mors, W.B., 1994. Furocoumarins from the rhizomes of *Dorstenia brasiliensis*. *Phytochemistry* 36, 221–223.
- Leete, E., 1983. Chemistry of the tropane alkaloids. 31. Biosynthesis of cocaine and cuscohygrine from [1-C-14]-labeled acetate and [4-H-3]-labeled phenylalanine in *Erythroxylon coca*. *Phytochemistry* 22, 699–704.
- Lewis, D.A., Fields, W.N., Shaw, G.P., 1999. A natural flavonoid present in unripe plantain banana pulp (*Musa sapientum* L. var. *paradisica*) protects the gastric mucosa from aspirin-induced erosions. *Journal of Ethnopharmacology* 65, 283–288.
- Lorenzi, H., 1992. *Árvores Brasileiras: Manual de Identificação de Plantas Arbóreas Nativas do Brasil*. Plantarum, Nova Odessa.
- Lorenzi, H., 1998. *Árvores Brasileiras: Manual de Identificação de Plantas Arbóreas Nativas do Brasil*. Plantarum, Nova Odessa.
- Lorenzi, H., Matos, F.J.A., 2002. *Plantas Medicinais no Brasil: Nativas e Exóticas*. Plantarum, Nova Odessa.
- Lorenzi, H., 2003. *Árvores Exóticas no Brasil*. Plantarum, Nova Odessa.
- Lorenzo, D., Loayza, I., Leigue, L., Frizzo, C., Dellacassa, E., Moyna, P., 2001. Asaricin, the main component of *Ocotea opifera* Mart. essential oil. *Natural Product Letters* 15, 163–170.
- Marder, M., Viola, H., Wasowski, C., Fernandez, S., Medina, J.H., Paladini, A.C., 2003. 6-Methylapigenin and hesperidin: new valeriana flavonoids with activity on the CNS. *Pharmacology Biochemistry and Behavior* 75, 537–545.
- Martius, K.F.P.V., 1939. *Natureza, Doenças, Medicina e Remédios dos Índios Brasileiros (1844)*. Companhia Editorial Nacional, Sao Paulo.
- Melo Cavalcante, A.A., Rubensam, G., Picada, J.N., Gomes Da Silva, E., Fonseca Moreira, J.C., Henriques, J.A., 2003. Mutagenicity, antioxidant potential and antimutagenic activity against hydrogen peroxide of cashew (*Anacardium occidentale*) apple juice and cajuina. *Environmental and Molecular Mutagenesis* 41, 360–369.
- Melzig, M.F., Putscher, I., Henklein, P., Haber, H., 2000. In vitro pharmacological activity of the tetrahydroisoquinoline salsolinol present in products from *Theobroma cacao* L. like cocoa and chocolate. *Journal of Ethnopharmacology* 73, 153–159.
- Ming, L.C., 1995. Levantamento de plantas medicinais na Reserva Extrativista “Chico Mendes”—ACRE. Doctorate Thesis, Instituto de Biociências da Universidade Estadual Paulista “Julio Mesquita Filho”—Campus Botucatu, 174 p.
- Missouri Botanical Garden, 2006. <http://mobot.mobot.org/W3T/Search/vast.html> (accessed 03 June, 2006).
- Moreau, R.A., Hicks, K.B., 2005. The composition of corn oil obtained by the alcohol extraction of ground corn. *Journal of the American Oil Chemists Society* 82, 809–815.
- Mota, M.L., Thomas, G., Barbosa Filho, J.M., 1985. Anti-inflammatory actions of tannins isolated from the bark of *Anacardium occidentale* L. *Journal of Ethnopharmacology* 13, 289–300.
- Nagase, H., Omae, N., Omori, A., Nakagawasai, O., Tadano, T., Yokosuka, A., Sashida, Y., Mimaki, Y., Yamakuni, T., Ohizumi, Y., 2005. Nobiletin and its related flavonoids with CRE-dependent transcription-stimulating and neuritegenic activities. *Biochemical and Biophysical Research Communications* 337, 1330–1336.
- Olatunji, L.A., Okwusidi, J.I., Soladoye, A.O., 2005. Antidiabetic effect of *Anacardium occidentale* stem-bark in fructose-diabetic rats. *Le Pharmacien Biologiste* 43, 589–593.
- Osman, H., Nasarudin, R., Lee, S.L., 2004. Extracts of cocoa (*Theobroma cacao* L.) leaves and their antioxidant potential. *Food Chemistry* 86, 41–46.
- Ozturk, N., Baser, K.H.C., Aydin, S., Ozturk, Y., Calis, I., 2002. Effects of *Gentiana lutea* ssp. *symphyandra* on the central nervous system in mice. *Phytotherapy Research* 16, 627–631.

- Pachter, I.J., Zacharias, D.E., Ribeiro, O., 1959. Indole alkaloids of acer-saccharinum (the silver maple), *Dictyoloma incanescens*, *Piptadenia colubrina*, and *Mimosa hostilis*. The Journal of Organic Chemistry 24, 1285–1287.
- Pannangpetch, P., Vuttivirojana, A., Kularbkaew, C., Tesana, S., Kongyingyoes, B., Kukongviriyapan, V., 2001. The antiulcerative effect of Thai *Musa* species in rats. Phytotherapy Research 15, 407–410.
- Pari, L., Maheswari, J.U., 1999. Hypoglycaemic effect of *Musa sapientum* L-in alloxan-induced diabetic rats. Journal of Ethnopharmacology 68, 321–325.
- Phillips, K.M., Ruggio, D.M., Ashraf-Khorassani, M., 2005. Phytosterol composition of nuts and seeds commonly consumed in the United States. Journal of Agricultural and Food Chemistry 53, 9436–9445.
- Pino, J.A., Ortega, A., Marbot, R., Aguero, J., 2003. Volatile components of banana fruit (*Musa sapientum* L.) “Indio” from Cuba. Journal of Essential Oil Research 15, 79–80.
- Pio Correa, M., 1926. Dicionário das Plantas Uteis do BRASIL e das Exóticas Cultivadas, vol. 6. Ministério da Agricultura, Rio de Janeiro.
- Pires, J.M., 1978. Plantas icotóxicas: aspecto da botânica sistemática. Anais do V Simposio Brasileiro de Plantas Mediciniais, 37.
- Plowman, T., 1979. The Identity of Amazonian and Trujillo Coca. Botanical Museum Leaflets, vol. 27. Harvard University, pp. 45–68. Pubmed, <http://www.ncbi.nlm.nih.gov/entrez/> (accessed August, 2005).
- Rahman, W., Ishratullah, K., Wagner, H., Seligmann, O., Chari, V.M., Osterdahl, B.G., 1978. Prunin-6-ortho-para-coumarate—new acylated flavanone glycoside from *Anacardium occidentale*. Phytochemistry 17, 1064–1065.
- Rahmat, A., Kumar, V., Fong, L.M., Endrini, S., Sani, H.A., 2004. Determination of total antioxidant activity in three types of local vegetables shoots and the cytotoxic effect on their ethanolic extracts against different cancer cell lines. Asia Pacific Journal of Clinical Nutrition 13, 308.
- Ramiro, E., Franch, A., Castellote, C., Perez-Cano, F., Permanyer, J., Izquierdo-Pulido, M., Castell, M., 2005. Flavonoids from *Theobroma cacao* down-regulate inflammatory mediators. Journal of Agricultural and Food Chemistry 53, 8506–8511.
- Rodrigues, E., 2005. A Parceria Universidade—empresa privada na produção de fitoterápicos no Brasil. Revista FÁRMACOS e Medicamentos 37, 30–39.
- Rodrigues, E., 2006. Plants and animals utilized as medicines in the Jau National Park (JNP), Brazilian Amazon. Phytotherapy Research 20, 378–391.
- Sanchez-Rababada, F., Jauregui, O., Casals, I., Andres-Lacueva, C., Izquierdo-Pulido, M., Lamuela-Raventos, R.M., 2003. Liquid chromatographic/electrospray ionization tandem mass spectrometric study of the phenolic composition of cocoa (*Theobroma cacao*). Journal of Mass Spectrometry 38, 35–42.
- Santos, R.V.H., Leôncio, M.P., Afatpour, P., 1994. Efeitos farmacológicos e ação diurética do extrato hidroalcoólico de *Selaginella convoluta* Spring. Anais do XIII Simposio Brasileiro de Plantas Mediciniais, 40.
- Santos, R.F., Galduroz, J.C.F., Barbieri, A., Castiglioni, M., Ytaya, L., Bueno, O.F.A., 2003. Cognitive Performance, SPECT, and Blood Viscosity in Elderly Non-demented People using Ginkgo biloba, vol. 36. Pharmacopsychiatry, Germany, pp. 1–8.
- Scapagnini, G., Butterfield, D.A., Colombrina, C., Sultana, R., Pascale, A., Calabrese, V., 2004. Ethyl ferulate, a lipophilic polyphenol, induces HO-1 and protects rat neurons against oxidative stress. Antioxidants and Redox Signaling 6, 811–818.
- Schewe, T., Kuhn, H., Sies, H., 2002. Flavonoids of cocoa inhibit recombinant human 5-lipoxygenase. The Journal of Nutrition 132, 1825–1829.
- Schreiber, L., Franke, R., Hartmann, K.D., Ranathunge, K., Steudle, E., 2005. The chemical composition of suberin in apoplastic barriers affects radial hydraulic conductivity differently in the roots of rice (*Oryza sativa* L. cv. IR64) and corn (*Zea mays* L. cv. Helix). Journal of Experimental Botany 56, 1427–1436.
- Schultes, R.E., 1969. Hallucinogens of plant origin. Science 163, 245–254.
- Schultes, R.E., Raffauf, R.F., 1990. The Healing Forest. Medicinal and Toxic Plants of the Northwest Amazonia, vol. 2. Discorides Press, Oregon.
- Silva, V.A., Andrade, L.H.C., 1998. Etnobotânica Xucuru: plantas medicinais. Revista Brasileira de Farmacologia 79, 33–36.
- Singh, Y.N., Dryden, W.F., 1985. Muscle paralyzing effect of the juice from the trunk of the banana tree. Toxicon 23, 973–981.
- Sousa, G.S., 1974. Notícia do Brasil (1587). Brasiliensia Documenta, Sao Paulo.
- Spix, J.B.V., Martius, K.F.P.V., 1976. Viagem Pelo Brasil (1817–1820). Editora Melhoramentos, Sao Paulo.
- Stark, T., Bareuther, S., Hofmann, T., 2005. Sensory-guided decomposition of roasted cocoa nibs (*Theobroma cacao*) and structure determination of taste-active polyphenols. Journal of Agricultural and Food Chemistry 53, 5407–5418.
- Thomas, M.B., 2001. An Analysis of the Pataxó Pharmacopeia of Bahia, Brazil, Using an Object Oriented Database Model. Doctorate Thesis. University of Florida.
- Torres, D.E.G., Assunção, D., Mancini, P., Torres, R.P., Mancini, J., 2002. Antioxidant activity of macambo (*Theobroma bicolor* L.) extracts. European Journal of Lipid Science and Technology 104, 278–281.
- Trevisan, M.T.S., Pfundstein, B., Haubner, R., Wurtele, G., Spiegelhalter, B., Bartsch, H., Owen, R.W., 2006. Characterization of alkyl phenols in cashew (*Anacardium occidentale*) products and assay of their antioxidant capacity. Food and Chemical Toxicology 44, 188–197.
- Turner, C.E., Ma, C.Y., Elsohly, M.A., 1981. Constituents of erythroxyton-coca. 2. Gas-chromatographic analysis of cocaine and other alkaloids in coca leaves. Journal of Ethnopharmacology 3, 293–298.
- Tyman, J.H.P., Tychopoulos, V., Chan, P., 1984. Long-chain phenols. 25. Quantitative-analysis of natural cashew nut-shell liquid (*Anacardium occidentale*) by high-performance liquid-chromatography. Journal of Chromatography 303, 137–150.
- Uchiyama, T., Hara, S., Makino, M., Fujimoto, Y., 2002. *Dorstenia brasiliensis*. Seco-Adianane-type triterpenoids from *Dorstenia brasiliensis* (Moraceae). Phytochemistry 60, 761–764.
- USPTO (United States Patent and Trademark Office), 2005. <http://www.uspto.gov> (accessed 14 August, 2005).
- Van den Berg, M.E., Silva, M.H.L., 1988. Contribuição ao Conhecimento da Flora Medicinal de Roraima. Supl. Acta Amazônica 18, 23–25.
- Veloso, J.M.C., 1976. Plantas Fluminenses Descritas por Frei Veloso (1741–1811). Separata de Anais da Biblioteca Nacional, Rio de Janeiro.
- Wagener, Z., 1964. “Zoobiblión” (Livro de Animais do Brasil) (1654). Brasiliensia Documenta, Sao Paulo.
- Web of Science. <http://scientific.thomson.com/products/wos/> (accessed June 2006).
- WHO (World Health Organization), [http://www.who.int/mental\\_health/en/](http://www.who.int/mental_health/en/) (accessed 14 May, 2006).
- Yan, Z.G., Wang, C.Z., 2006. Wound-induced green leaf volatiles cause the release of acetylated derivatives and a terpenoid in maize. Phytochemistry 67, 34–42.
- Yin, L.L., Zhu, X.Z., 2005. The involvement of central cholinergic system in (+)-matrine-induced antinociception in mice. Pharmacology Biochemistry and Behavior 80, 419–425.
- Zamble, A., Carpentier, M., Kandoussi, A., Sahpaz, S., Petrault, O., Ouk, T., Hennuyer, N., Fruchart, J.C., Staels, B., Bordet, R., Duriez, P., Bailleul, F., Martin-Nizard, F., 2006. *Paullinia pinnata* extracts rich in polyphenols promote vascular relaxation via endothelium-dependent mechanisms. Journal of Cardiovascular Pharmacology 47, 599–608.